

(i) Results of tests and assays on the batch for potency, moisture, pH, residue on ignition, identity, and crystallinity.

(ii) Samples required: 10 packages, each containing approximately 500 milligrams.

(b) *Tests and methods of assay*—(1) *Potency*. Proceed as directed in § 436.105 of this chapter, preparing the sample for assay as follows: Dissolve an accurately weighed sample in sufficient methyl alcohol to give a concentration of 1 milligram of erythromycin base per milliliter (estimated). Further dilute with 0.1M potassium phosphate buffer, pH 8.0 (solution 3), to the reference concentration of 1.0 microgram of erythromycin base per milliliter (estimated).

(2) [Reserved]

(3) *Moisture*. Proceed as directed in § 436.201 of this chapter.

(4) *pH*. Proceed as directed in § 436.202 of this chapter, using a 1.0 percent suspension in water.

(5) *Residue on ignition*. Proceed as directed in § 436.207(a) of this chapter.

(6) *Identity*. Proceed as directed in § 436.211 of this chapter, using the sample prepared as described in paragraph (b)(3) of that section.

(7) *Crystallinity*. Proceed as directed in § 436.203(a) of this chapter.

[39 FR 19149, May 30, 1974, as amended at 50 FR 19920, May 13, 1985]

**§ 452.25a Sterile erythromycin ethylsuccinate.**

(a) *Requirements for certification*—(1) *Standards of identity, strength, quality, and purity*. Erythromycin ethylsuccinate is the white, odorless, ethylsuccinate ester of erythromycin. It is so purified and dried that:

(i) It contains not less than 765 micrograms of erythromycin per milligram, calculated on an anhydrous basis.

(ii) It is sterile.

(iii) [Reserved]

(iv) Its moisture content is not more than 3.0 percent.

(v) Its pH is not less than 6.0 and not more than 8.5.

(vi) Its residue on ignition is not more than 1.0 percent.

(vii) It gives a positive identity test for erythromycin ethylsuccinate.

(viii) It is crystalline.

(2) *Labeling*. It shall be labeled in accordance with the requirements of § 432.5(b) of this chapter.

(3) *Requests for certification; samples*. In addition to the requirements of § 431.1 of this chapter, each such request shall contain:

(i) Results of tests and assays on the batch for potency, sterility, moisture, pH, residue on ignition, identity, and crystallinity.

(ii) Samples required:

(a) For all tests except sterility: 10 packages, each containing approximately 500 milligrams.

(b) For sterility testing: 20 packages, each containing approximately 600 milligrams.

(b) *Tests and methods of assay*—(1) *Potency*. Proceed as directed in § 436.105 of this chapter, preparing the sample for assay as follows: Dissolve an accurately weighed sample in sufficient methyl alcohol to give a concentration of 1 milligram of erythromycin base per milliliter (estimated). Further dilute with 0.1M potassium phosphate buffer, pH 8.0 (solution 3), to the reference concentration of 1.0 microgram of erythromycin base per milliliter (estimated).

(2) *Sterility*. Proceed as directed in § 436.20 of this chapter, using the method described in paragraph (e)(2) of that section.

(3) [Reserved]

(4) *Moisture*. Proceed as directed in § 436.201 of this chapter.

(5) *pH*. Proceed as directed in § 436.202 of this chapter, using a 1.0 percent suspension in water.

(6) *Residue on ignition*. Proceed as directed in § 436.207(a) of this chapter.

(7) *Identity*. Proceed as directed in § 436.211 of this chapter, using the sample preparation method described in paragraph (b)(3) of that section.

(8) *Crystallinity*. Proceed as directed in § 436.203(a) of this chapter.

[39 FR 19149, May 30, 1974, as amended at 50 FR 19920, May 13, 1985]

**§ 452.30a Sterile erythromycin gluceptate.**

(a) *Requirements for certification*—(1) *Standards of identity, strength, quality, and purity*. Erythromycin gluceptate is the white powder of the glucoheptonic

acid salt of erythromycin or a mixture of two or more such salts. It is freely soluble in water, alcohol, and methyl alcohol. It is slightly soluble in acetone and chloroform, but is practically insoluble in ether. It is so purified and dried that:

(i) It contains not less than 600 micrograms of erythromycin per milligram, calculated on an anhydrous basis. If it is packaged for dispensing, its potency is satisfactory if it is not less than 90 percent and not more than 115 percent of the number of milligrams of erythromycin that it is represented to contain.

(ii) It is sterile.

(iii) [Reserved]

(iv) It is nonpyrogenic.

(v) Its moisture content is not more than 5.0 percent.

(vi) Its pH in an aqueous solution containing 25 milligrams per milliliter is not less than 6.0 nor more than 8.0.

(vii) It gives a positive identity test for erythromycin gluceptate.

(2) [Reserved]

(3) *Labeling.* It shall be labeled in accordance with the requirements of § 432.5 of this chapter.

(4) *Requests for certification; samples.* In addition to the requirements of § 431.1 of this chapter, each such request shall contain:

(i) Results of tests and assays on the batch for potency, sterility, pyrogens, moisture, pH, and identity.

(ii) Samples required:

(a) If the batch is packaged for repackaging or for use as an ingredient in the manufacture of another drug:

(1) For all tests except sterility: 10 packages, each containing not less than 300 milligrams.

(2) For sterility testing: 20 packages, each containing approximately 300 milligrams.

(b) If the batch is packaged for dispensing:

(1) For all tests except sterility: A minimum of 12 immediate containers of the batch.

(2) For sterility testing: 20 immediate containers, collected at regular intervals throughout each filling operation.

(b) *Tests and methods of assay*—(1) *Potency.* Proceed as directed in § 436.105 of this chapter, preparing the sample for assay as follows: If the batch is pack-

aged for repackaging or for use in manufacturing another drug, dissolve an accurately weighed sample in sufficient methyl alcohol to give a concentration of 10 milligrams of erythromycin base per milliliter (estimated). Dilute this solution further with sufficient 0.1M potassium phosphate buffer, pH 8.0 (solution 3), to give a stock solution containing 1.0 milligram of erythromycin base per milliliter (estimated). If it is packaged for dispensing, reconstitute as directed in the labeling. Then using a suitable hypodermic needle and syringe, remove all of the withdrawable contents if it is represented as a single dose container; or if the labeling specifies the amount of potency in a given volume of the resultant preparation, remove an accurately measured representative portion from each container. Dilute with solution 3 to give a stock solution of convenient concentration. Further dilute the stock solution with solution 3 to the reference concentration of 1.0 microgram of erythromycin base per milliliter (estimated).

(2) *Sterility.* Proceed as directed in § 436.20 of this chapter, using the method described in paragraph (e)(1) of that section.

(3) *Pyrogens.* Proceed as directed in § 436.32(b) of this chapter, using a solution containing 30 milligrams of erythromycin per milliliter.

(4) [Reserved]

(5) *Moisture.* Proceed as directed in § 436.201 of this chapter.

(6) *pH.* Proceed as directed in § 436.202 of this chapter, using a concentration of 25 milligrams per milliliter.

(7) *Identity.* Proceed as directed in § 436.211 of this chapter, using the sample preparation method described in paragraph (b)(2) of that section.

[39 FR 19149, May 30, 1974, as amended at 46 FR 16685, Mar. 13, 1981; 50 FR 19920, 19921, May 13, 1985]

#### **§ 452.32a Sterile erythromycin lactobionate.**

(a) *Requirements for certification*—(1) *Standards of identity, strength, quality, and purity.* Erythromycin lactobionate is the white to off-white powder of the lactobionate salt of erythromycin or a mixture of two or more such salts. It is so purified and dried that: